

Appendix C. Climate Overview for the Oak Ridge Area

Appendix C. Climate Overview for the Oak Ridge Area

C. 1 Regional Climate

The climate of the Oak Ridge area and its surroundings may be broadly classified as humid subtropical. The term “humid” indicates that the region receives an overall surplus of precipitation compared to the level of evapotranspiration that is normally experienced throughout the year. The “subtropical” nature of the local climate indicates that the region experiences warm to hot summers and cool winters. Such areas typically experience significant changes in temperature between summer and winter.

Local winters are characterized by synoptic weather systems that often produce significant precipitation events every 3 to 5 days. These wet periods are occasionally followed by arctic air outbreaks. Although snow and ice are not associated with many of these systems, occasional snowfall does occur. Winter cloud cover tends to be enhanced by the regional terrain (cold air wedging).

Severe thunderstorms are the most frequent during spring but can occur at any time during the year. The Cumberland Mountains and the Cumberland Plateau often inhibit the intensity of severe systems that traverse the region (due to the downward momentum created as the storms move off of the higher terrain). Summers are characterized by very warm, humid conditions. Occasional frontal systems may produce organized lines of thunderstorms (and rare damaging tornados). More frequently, however, summer precipitation results from “air mass” thundershowers that form as a consequence of daytime heating, rising humid air, and local terrain features. Although adequate precipitation usually occurs during the fall, the months of August through October represent the driest period of the year. The occurrence of precipitation during the fall tends to be less cyclic than during other seasons and is occasionally enhanced by decaying tropical systems moving north from the Gulf of Mexico. During November, winter-type cyclones again begin to dominate the weather and continue to do so until May.

Decadal-scale climate change has recently affected the East Tennessee region. Some of these changes are related to the hemispheric effects caused by the El Niño–Southern Oscillation (ENSO), the Pacific Decadal Oscillation (PDO), and the Atlantic-Multidecadal Oscillation (AMO). The first two patterns, having cycles of 3 to 7 years and about 40 years, respectively, affect Pacific Ocean sea surface temperatures. The AMO affects Atlantic sea surface temperature (again, having a cycle of about 40 years). All of these patterns can collectively modulate regional temperature and precipitation trends with respect to East Tennessee. The AMO has recently shifted from a cold to warm sea surface temperature phase (mid-1990s) while the PDO appears to be entering a cool sea surface temperature phase (since 2000). Also, the ENSO pattern has more frequently brought about warmer Eastern Pacific sea surface temperatures. Additionally, there is some evidence that human-induced climate change may be affecting the area as well (about +1.7°C in Oak Ridge during the last 30 years). The recent warming appears to have lengthened the growing season (i.e., the period with temperatures above 0°C) by about 2 to 3 weeks over the last 30 years.

C.2 Winds

Five major terrain-related wind regimes regularly affect the Great Valley of Eastern Tennessee: pressure-driven channeling, downward-momentum transport or vertically coupled flow, forced channeling, along-valley thermal circulations, and mountain-valley circulations. Pressure-driven channeling and vertically coupled flow (unstably stratified conditions) affect wind flow on scales comparable to that of the Great Valley (hundreds of kilometers). Forced channeling occurs on similar scales but is also quite important at smaller spatial scales, such as that of the local ridge-and-valley (Birdwell 1996). Along-valley and mountain-valley circulations are thermally driven and occur within a large range of spatial scales. Thermal flows are more prevalent under conditions of clear skies and low humidity.

Pressure-driven channeling, in its simplest essence, is the redirection of synoptically induced wind flow through a valley channel. The direction of wind flow through the valley is determined by the pressure gradient superimposed on the valley's axis (Whiteman 2000). The process is affected by Coriolis forces, a leftward deflection of winds (in the Northern Hemisphere). Eckman (1998) suggested that pressure-driven channeling plays a significant role in the Great Valley. Winds driven purely by such a process shift from up-valley to down-valley flow or conversely as "weather"-induced flow shifts across the axis of the Great Valley. Since the processes involved in pressure-driven flow primarily affect the horizontal motion of air, the presence of a temperature inversion enhances flow significantly. Weak vertical air motion and momentum associated with such inversions allow different layers of air to slide over each other (Monti et al. 2002).

Forced channeling is defined as the direct deflection of wind by terrain. This form of channeling necessitates some degree of vertical motion transfer, implying that the mechanism is less pronounced during temperature-inversion conditions. Although forced channeling may result from interactions between large valleys and mountain ranges (such as the Great Valley and the surrounding mountains), the mechanism is especially important in narrow, small valleys such as those on the Oak Ridge Reservation (Kossman and Sturman 2002).

Large-scale forced channeling occurs regularly within the Great Valley when northwest to north winds (perpendicular to the axis of the central Great Valley) coincide with vertically coupled flow. The phenomenon sometimes results in a split flow pattern (winds southwest of Knoxville moving down-valley and those to the east of Knoxville moving up-valley). The causes of such a flow pattern may include the shape characteristics of the Great Valley (Kossman and Sturman 2002) but also may be related to the specific location of the Cumberland and Smoky Mountains relative to upper level wind flow (Eckman 1998). The convex shape of the Great Valley with respect to a northwest wind flow may lead to a divergent wind flow pattern in the Knoxville area. This results in downward air motion. Additionally, horizontal flow is reduced by the windward mountain range (Cumberland Mountains), which increases buoyancy and Coriolis effects (Froude and Rossby ratios in the meteorological field). Consequently, the leeward mountain range (Smoky Mountains) becomes more effective at blocking or redirecting the winds.

Vertically coupled winds occur when the atmosphere is unstable (characterized by cooler temperatures aloft). When a strong horizontal wind component is also present (as in conditions behind a winter cold front), winds "ignore" the terrain, flowing over it in roughly in the same direction as the winds aloft. This phenomenon is a consequence of the horizontal transport and momentum aloft being transferred to the surface. However, Coriolis effects may turn the winds by up to 25° to the left (Birdwell 1996).

Thermally driven winds are common in areas of significantly complex terrain. These winds occur as a result of pressure and temperature differences caused by varied surface-air energy exchange at similar altitudes along a valley's axis, sidewalls, and/or slopes. Thermal flows operate most effectively when synoptic winds are light and when thermal differences are exacerbated by clear skies and low humidity (Whiteman 2000). Ridge-and-valley terrain may be responsible for enhancing or inhibiting such air flow, depending on the ambient weather conditions. Eckman (1998) suggested that the presence of daytime up-valley winds and night time down-valley (drainage) flows between the ridge-and-valley terrain of the Oak Ridge area tended to reverse at about 9:00 to 11:00 a.m. and at about 5:00 to 7:00 p.m. local time, respectively. The terrain-following nature of drainage winds suggests that they would be more directly impacted by the presence of the ridge-and-valley than daytime flows, which tend to be accompanied by significant vertical motions.

Figures C.1 thru C.17 display wind roses for each of the eight Oak Ridge Reservation meteorological towers during 2007 (Towers MT1, MT2, MT3, MT4, MT6, MT7, MT9, and MT10). The wind roses represent typical trends and should be used with caution.

A wind rose depicts the typical distribution of wind speed and direction for a given location. The winds are represented in terms of the direction from which they originate. The rays emanating from the center correspond to points of the compass. The length of each ray is related to the frequency that winds




Fig. C.2. Wind rose for ETP Meteorological Tower 1 for data taken at 60 m above ground level, 2007.

January and February average temperatures have seen increases of 7.0°F and 4.4°F, respectively. This dramatic increase is probably due to the fact that the Arctic has seen the largest increase in temperatures of anywhere in the Northern Hemisphere over the last 30 years. During the months of January and February, much of the air entering eastern Tennessee comes from the Arctic. As a result, Oak Ridge temperatures have warmed more dramatically during these months. Early spring temperatures (March–April) have risen by about 3°F whereas most months in the remainder of the calendar year saw temperature increases of about 2.5°F. The smallest increases, however, are observed for June and July. Average temperatures increased by less than 2°F during those months and were entirely the result of minimum temperature increases. Overall, annual minimum temperatures seem to have increased more dramatically (by 4.1°F) than maximum temperatures (by 2.1°F). Of the summer months, August and September increased the most dramatically (by about 2°F). For the most recent decade (1998–2007), August average temperatures are now about equal to those of July.

Decadal precipitation averages suggest some important changes in precipitation patterns in Oak Ridge over the period of 1978 to 2007. Although overall precipitation has remained within a window of about 48 to 56 in. annually, there have been some recent decadal shifts in the patterns of rainfall on a monthly or seasonal scale. In particular, precipitation has tended to increase during the late winter and early spring (February through April) by about 1 in./month. Conversely, the late summer and early fall months (August through October) have seen slight decreases in precipitation (averaging about 0.50 in./month). Overall, annual precipitation averaged an increase of 4 in. during the 1998–2007 decade vs the




Fig. C.3. Wind rose for ETPP Meteorological Tower 7 for data taken at 10 m above ground level, 2007.

1978–1987 period (see also Table C.2). Despite this average increase in rainfall, 2007 was the driest year on record in Oak Ridge (35.87 in.). This statistic encompasses the period from 1948 to 2007. Next to 2007, the second driest year on record occurred in 1958 (37.43 in.).

The previously discussed increase in winter temperatures has apparently affected monthly and annual snowfall amounts as well. During the decade of 1978 to 1987, snowfall averaged about 13 in. annually in Oak Ridge. However, during the most recent decade (1998–2007), snowfall has averaged only 4 in. This decrease seems to have occurred largely since the mid-1990s.

Figures C.18 thru C.21 provide typical wind roses for Tower MT2 (“C”) during light, moderate, heavy, and all precipitation events, respectively. The precipitation classes are defined by the National Weather Service as follows:

- light: trace to 0.10 in./h
- moderate: 0.11 to 0.30 in./h
- heavy: more than 0.30 in./h

The meteorological record from ORNL’s Tower C was used because it is centrally located within the Oak Ridge Reservation.




Fig. C.4. Wind rose for ETPP Meteorological Tower 7 for data taken at 30 m above ground level, 2007.

Hourly values of subfreezing temperatures in Oak Ridge are presented in Table C.3 for the years 1985 to 2007. During the mid-to-late 1980s, a typical year experienced about 900 to 1000 h of subfreezing temperatures. In recent years, the value has fallen to approximately 600 h.

C.4 Stability

The local ridge-and-valley terrain plays a role in the development of stable surface air under certain conditions and influences the dynamics of air flow. Although ridge-and-valley terrain creates identifiable patterns of association during unstable conditions as well, strong vertical mixing and momentum tend to significantly reduce these effects. “Stability” describes the tendency of the atmosphere to mix or overturn. Consequently, dispersion parameters are influenced by the stability characteristics of the atmosphere. Stability classes range from “A” (very unstable) to “G” (very stable). The “D” stability class represents a neutral state.

The suppression of vertical motions during stable conditions increases the frequency with which air motion is impacted by the local terrain. Conversely, stable conditions isolate wind flows within the ridge-and-valley terrain from the effects of more distant terrain features and from winds aloft. These effects are particularly true with respect to mountain waves. Deep stable layers of air tend to reduce the vertical space available for oscillating vertical air motions caused by local mountain ranges (Smith et al. 2002). This effect on mountain wave formation may be important with regard the impact that the nearby Cumberland Mountains may have on local air flow.




Fig. C.5. Wind rose for ORNL Meteorological Tower 2 for data taken at 10 m above ground level, 2007.

A second factor that may decouple large-scale wind flow effects from local ones (and thus produce stable surface layers) occurs with overcast sky conditions. Clouds overlying the Great Valley may warm due to direct insolation on the cloud tops. Warming may also occur within the clouds as latent energy, which is released due to the condensation of moisture. Surface air underlying the clouds may remain relatively cool (as it is cut off from direct exposure to the sun). Consequently, the vertical temperature gradient associated with the air mass becomes more stable (Lewellen and Lewellen 2002). Long wave cooling of fog decks has also been observed to help modify stability in the surface layer (Whiteman et al. 2001).

Stable boundary layers typically form as a result of radiational cooling processes near the ground (Van De Weil et al. 2002); however, they are also influenced by the mechanical energy supplied by horizontal wind motion (which is in turn influenced by the large-scale “weather”-related pressure gradient). Ridge-and-valley terrain may have a significant ability to block such winds and their associated mechanical energy (Carlson and Stull 1986). Consequently, enhanced radiational cooling at the surface results since there is less wind energy available to remove chilled air.

Stable boundary layers also exhibit intermittent turbulence that has been associated with a number of the above factors. The process results from a “give-and-take” between the effects of friction and radiational cooling. As a stable surface layer intensifies via a radiation cooling process, it tends to decouple from air aloft, thereby reducing the effects of surface friction. The upper air layer responds with




Fig. C.6. Wind rose for ORNL Meteorological Tower 2 for data taken at 30 m above ground level, 2007.

an acceleration in wind speed. Increased wind speed aloft results in an increase in mechanical turbulence and wind shear at the boundary with the stable surface layer. Eventually, the turbulence works into the surface layer and weakens it. As the inversion weakens, friction again increases, reducing winds aloft. The reduced wind speeds aloft allow enhanced radiation cooling at the surface, which reintensifies the inversion and allows the process to start again. Van De Weil et al. (2002) have shown that cyclical temperature oscillations up to 4°C may result from these processes. Since these intermittent processes are driven primarily by large-scale horizontal wind flow and radiational cooling of the surface, ridge-and-valley terrain significantly affects these oscillations.

Figures C.22 thru C.28 provide wind roses for ORNL Tower MT2 at 30 m with respect to Stability A thru G during 2007. Stability “A” (unstable) conditions show a strong preference for winds from the south half of the compass. Stability “D” conditions (neutral), which also tend to correspond to higher wind speed, show a significant preference for winds from the west and west-northwest. During very stable conditions (F and G stability), winds shows a preference for eastnortheast directions (likely down valley “cold air” drainage flow).




Fig. C.7. Wind rose for ORNL Meteorological Tower 2 for data taken at 100 m above ground level, 2007.




Fig. C.9. Wind rose for ORNL Meteorological Tower 3 for data taken at 30 m above ground level, 2007.




Fig. C.10. Wind rose for ORNL Meteorological Tower 4 for data taken at 10 m above ground level, 2007.




Fig. C.11. Wind rose for ORNL Meteorological Tower 4 for data taken at 30 m above ground level, 2007.




Fig. C.12. Wind rose for ORNL Meteorological Tower 10 for data taken at 10 m above ground level, 2007.




Fig. C.14. Wind rose for ORNL Meteorological Tower 6 for data taken at 30 m above ground level, 2007.




Fig. C.16. Wind rose for Y-12 Meteorological Tower 9 for data taken at 15 m above ground level, 2007.




Fig. C.18. Wind rose for ORNL Meteorological Tower 2 for data taken at 30 m above ground level during light precipitation events (trace-0.10 in.), 1998-2007.




Fig. C.19. Wind rose for ORNL Meteorological Tower 2 for data taken at 30 m above ground level during moderate precipitation events (0.10–0.30 in.), 1998–2007.




Fig. C.20. Wind rose for ORNL Meteorological Tower 2 for data taken at 30 m above ground level during heavy precipitation events (> 0.30 in.), 1998–2007.




Fig. C.21. Wind rose for ORNL Meteorological Tower 2 for data taken at 30 m above ground level during all precipitation events, 1998–2007.




Fig. C.22. Wind rose for ORNL Meteorological Tower 2, stability class A, for data taken at 30 m above ground level, 2007.




Fig. C.23. Wind rose for ORNL Meteorological Tower 2, stability class B, for data taken at 30 m above ground level, 2007.




Fig. C.24. Wind rose for ORNL Meteorological Tower 2, stability class C, for data taken at 30 m above ground level, 2007.




Fig. C.25. Wind rose for ORNL Meteorological Tower 2, stability class D, for data taken at 30 m above ground level, 2007.




Fig. C.26. Wind rose for ORNL Meteorological Tower 2, stability class E, for data taken at 30 m above ground level, 2007.




Fig. C.27. Wind rose for ORNL Meteorological Tower 2, stability class F, for data taken at 30 m above ground level, 2007.




Fig. C.28. Wind rose for ORNL Meteorological Tower 2, stability class G, for data taken at 30 m above ground level, 2007.

Table C.1. Climate normals (1978–2007) and extremes (1948–2007) for Oak Ridge, Tennessee (Town Site), with 2007 comparisons

Monthly variables	January	February	March	April	May	June	July	August	September	October	November	December	Annual
Temperature, °C (°F)													
30-year average max	8.1 (46.5)	10.8 (51.5)	16.4 (61.5)	21.6 (70.8)	25.8 (78.5)	29.6 (85.3)	31.4 (88.6)	31.2 (88.1)	27.7 (81.9)	21.9 (71.5)	15.5 (59.9)	9.7 (49.4)	20.8 (69.5)
2007 average max	10.2 (50.4)	8.6 (47.5)	20.6 (69.0)	19.8 (67.6)	28.0 (82.4)	31.2 (88.1)	30.5 (86.9)	35.3 (95.3)	30.4 (86.7)	24.3 (75.8)	15.6 (60.1)	12.6 (54.6)	22.3 (72.1)
60-year record max	25 (77)	26 (79)	30 (86)	33 (92)	34 (93)	38 (101)	41 (105)	39 (103)	39 (102)	32 (90)	28 (83)	26 (78)	41 (105)
30-year average min	-2.4 (27.7)	-1.0 (30.2)	2.8 (37.0)	7.2 (45.0)	12.3 (54.2)	17.0 (62.6)	19.6 (67.3)	19.1 (66.3)	15.1 (59.2)	8.2 (46.7)	2.9 (37.3)	-1.0 (30.2)	8.3 (47.0)
2007 average min	-0.3 (31.5)	-2.3 (27.9)	7.2 (45.0)	7.0 (44.6)	13.8 (56.9)	17.7 (63.9)	17.6 (63.7)	22.1 (71.7)	16.7 (62.1)	11.4 (52.5)	3.1 (37.6)	2.2 (35.9)	9.8 (49.7)
60-year record min	-27 (-17)	-25 (-13)	-17 (1)	-7 (20)	-1 (30)	4 (39)	9 (49)	10 (50)	1 (33)	-6 (21)	-18 (0)	-22 (-7)	-27 (-17)
30-year average	2.8 (37.1)	4.9 (40.9)	9.6 (49.3)	14.4 (58.0)	19.1 (66.4)	23.3 (74.0)	25.5 (77.9)	25.1 (77.2)	21.4 (70.6)	15.1 (59.1)	9.2 (48.6)	4.3 (39.8)	14.6 (58.2)
2007 average	5.0 (41.0)	3.2 (37.7)	13.9 (57.0)	13.4 (56.1)	20.9 (69.7)	24.4 (76.0)	25.1 (77.1)	28.7 (83.7)	23.6 (74.4)	17.9 (64.2)	9.4 (48.9)	7.5 (45.5)	16.1 (60.9)
2007 dep from average	2.2 (3.9)	-1.8 (-3.2)	4.3 (7.7)	-1.1 (-1.9)	1.8 (3.3)	1.1 (2.0)	-0.4 (-0.8)	3.6 (6.5)	2.1 (3.8)	2.8 (5.1)	0.2 (0.3)	3.2 (5.7)	1.5 (2.7)
30-year average heating degree days, °C (°F)^a													
	477 (858)	376 (676)	269 (484)	129 (232)	38 (68)	2 (4)	0	0	14 (26)	114 (206)	271 (487)	430 (774)	2119 (3815)
30-year average cooling degree days, °C (°F)^a													
	0	0	2 (4)	14 (26)	65 (117)	156 (280)	226 (407)	215 (387)	111 (199)	17 (31)	1 (2)	0	807 (1453)
Precipitation, mm (in.)													
30-year average	122.5 (4.82)	119.7 (4.71)	126.0 (4.96)	114.3 (4.50)	118.1 (4.65)	108.0 (4.25)	137.2 (5.40)	82.8 (3.26)	95.5 (3.76)	69.1 (2.72)	122.2 (4.81)	124.8 (4.91)	1340 (52.75)
2007	74.4 (2.93)	33.3 (1.31)	83.6 (3.29)	134.9 (5.31)	69.1 (2.72)	40.4 (1.59)	106.5 (4.19)	18.0 (0.71)	62.8 (2.47)	70.6 (2.78)	128.8 (5.07)	88.9 (3.50)	911.4 (35.87)
2007 dep from average	-48.0 (-1.89)	-86.4 (-3.40)	-42.4 (-1.67)	20.6 (0.81)	-49.0 (-1.93)	-67.6 (-2.66)	-30.7 (-1.21)	-64.8 (2.55)	-32.8 (-1.29)	1.5 (0.06)	6.6 (0.26)	-35.8 (-1.41)	-429 (-16.88)
60-year max monthly	337.2 (13.27)	324.7 (12.78)	311.0 (12.24)	356.5 (14.03)	271.9 (10.70)	283.0 (11.14)	489.6 (19.27)	265.8 (10.46)	176.6 (6.95)	176.6 (6.95)	310.5 (12.22)	321.2 (12.64)	1939 (76.33)
60-year max 24-hr	108.0 (4.25)	131.6 (5.18)	120.4 (4.74)	158.5 (6.24)	112.0 (4.41)	94.0 (3.70)	124.8 (4.91)	190.1 (7.48)	129.8 (5.11)	67.6 (2.66)	130.1 (5.12)	130.1 (5.12)	190.1 (7.48)
60-year min monthly	23.6 (0.93)	21.3 (0.84)	54.1 (2.13)	22.4 (0.88)	20.3 (0.80)	13.5 (0.53)	31.3 (1.23)	13.7 (0.54)	Trace	Trace	34.8 (1.37)	17.0 (0.67)	911.4 (35.87)
Snowfall, mm (in.)													
30-year average	96.6 (3.8)	91.5 (3.6)	38.1 (1.5)	7.6 (0.3)	0	0	0	0	0	0	Trace	43.2 (1.7)	276.9 (10.9)
2007 totals	Trace	25.4 (1.0)	Trace	Trace	0	0	0	0	0	0	Trace	Trace	25.4 (1.0)
60-year max monthly	243.9 (9.6)	437.0 (17.2)	533.6 (21.0)	149.9 (5.9)	Trace	0	0	0	0	Trace	165.2 (6.5)	533.6 (21.0)	1052 (41.4)
60-year max 24-hr	210.9 (8.3)	287.1 (11.3)	304.9 (12.0)	137.2 (5.4)	Trace	0	0	0	0	Trace	165.2 (6.5)	304.9 (12.0)	304.9 (12.0)
Days w/Temp													
30-year max ≥ 32°C	0	0	0	0.1	1.0	5.5	14.7	12.9	4.2	0	0	0	38.4
2007 max ≥ 32°C	0	0	0	0	0	10	9	30	13	1	0	0	63
30-year min ≤ 0°C	21.9	16.4	11.7	2.7	0.1	0	0	0	0	2.0	11.0	19.6	85.4
2007 min ≤ 0°C	17	20	5	5	0	0	0	0	0	0	8	12	67
30-year max ≤ 0°C	2.9	1.3	0.2	0	0	0	0	0	0	0	0.1	1.6	6.1
2007 max ≤ 0°C	0	1	0	0	0	0	0	0	0	0	0	0	1
Days w/Precip													
30-year avg ≥ 0.01 in.	11.5	10.7	11.6	10.4	11.6	11.2	12.5	9.7	9.1	8.1	9.7	11.0	127.1
2007 days ≥ 0.01 in.	6	7	6	10	7	9	13	5	4	9	8	13	97
30-year avg ≥ 1.00 in.	1.3	1.2	1.5	0.8	1.4	1.3	1.4	0.7	1.2	0.7	1.3	1.3	14.1
2007 days ≥ 1.00 in.	1	0	1	2	0	0	1	0	1	1	1	1	9

Table C.2. Decadal climate change (1978–2007) for Oak Ridge, Tennessee (Town Site) with 2007 comparisons

Monthly variables	January	February	March	April	May	June	July	August	September	October	November	December	Annual
Temperature, °C (°F)													
1978-1987 Avg Max	5.7 (42.2)	9.2 (48.6)	15.6 (60.0)	20.9 (69.7)	25.4 (77.8)	29.7 (85.5)	31.3 (88.4)	30.6 (87.0)	27.3 (81.1)	21.5 (70.7)	15.3 (59.6)	9.2 (48.6)	20.1 (68.3)
1988-1997 Avg Max	8.9 (48.0)	11.8 (53.2)	16.9 (62.4)	21.9 (71.5)	25.7 (78.3)	29.5 (85.1)	31.8 (89.2)	31.2 (88.1)	27.6 (81.7)	22.0 (71.6)	14.8 (58.6)	9.5 (49.1)	21.0 (69.7)
1998-2007 Avg Max	9.6 (49.2)	11.4 (52.5)	16.7 (62.1)	21.8 (71.2)	26.3 (79.3)	29.7 (85.4)	31.2 (88.1)	31.8 (89.2)	28.3 (82.9)	22.4 (72.3)	16.4 (61.6)	10.3 (50.5)	21.3 (70.4)
Temp Change (1978-1987 vs. 1998-2007)	3.9 (7.0)	2.2 (3.9)	1.2 (2.1)	0.8 (1.5)	0.8 (1.5)	-0.1 (-0.1)	-0.2 (-0.3)	1.2 (2.2)	1.0 (1.8)	0.9 (1.6)	1.1 (2.0)	1.1 (1.9)	1.2 (2.1)
2007 Avg Max	10.2 (50.4)	8.6 (47.5)	20.6 (69.0)	19.8 (67.6)	28.0 (82.4)	31.2 (88.1)	30.5 (86.9)	35.3 (95.6)	30.4 (86.7)	24.3 (75.8)	15.6 (60.1)	12.6 (54.6)	22.3 (72.1)
1978-1987 Avg Min	-4.7 (23.5)	-2.6 (27.4)	1.4 (34.5)	5.9 (42.7)	11.5 (52.7)	15.9 (60.6)	18.7 (65.6)	18.1 (64.6)	14.3 (57.7)	7.3 (45.2)	2.9 (37.2)	-1.8 (28.7)	7.2 (45.0)
1988-1997 Avg Min	-1.7 (28.9)	-0.8 (30.6)	2.8 (37.1)	6.8 (44.2)	11.8 (53.3)	17.2 (62.9)	19.7 (67.4)	18.9 (66.1)	15.4 (59.7)	7.6 (45.7)	1.8 (35.2)	-1.0 (30.2)	8.2 (46.8)
1998-2007 Avg Min	-0.7 (30.7)	0.3 (32.6)	4.2 (39.5)	8.9 (48.1)	13.7 (56.6)	17.9 (64.3)	20.4 (68.8)	20.1 (68.2)	15.7 (60.2)	9.5 (49.1)	4.2 (39.5)	-0.2 (31.7)	9.5 (49.1)
Temp Change (1978-1987 vs. 1998-2007)	4.0 (7.2)	2.9 (5.2)	2.8 (5.0)	3.0 (5.4)	2.2 (3.9)	2.1 (3.7)	1.8 (3.2)	2.0 (3.6)	1.4 (2.5)	2.2 (3.9)	1.3 (2.3)	1.7 (3.0)	2.3 (4.1)
2007 Avg Min	-0.3 (31.5)	-2.3 (27.9)	7.2 (45.0)	7.0 (44.6)	13.8 (56.9)	17.7 (63.9)	19.6 (67.3)	22.1 (71.7)	16.7 (62.1)	11.4 (52.5)	3.1 (37.6)	2.2 (35.9)	9.9 (49.7)
1978-1987 Avg	0.5 (32.9)	3.4 (38.1)	8.5 (47.3)	13.4 (56.2)	18.5 (65.3)	22.8 (73.1)	25.0 (77.0)	24.3 (75.8)	20.9 (69.9)	14.4 (58.0)	9.1 (48.4)	3.7 (38.6)	13.7 (56.7)
1988-1997 Avg	3.6 (38.5)	5.5 (41.9)	9.9 (49.9)	14.3 (57.8)	18.8 (65.8)	23.4 (74.1)	25.7 (78.3)	25.1 (77.1)	21.5 (70.7)	14.8 (58.7)	8.3 (46.9)	4.3 (39.7)	14.6 (58.3)
1998-2007 Avg	4.4 (39.9)	5.8 (42.5)	10.4 (50.8)	15.5 (59.9)	20.0 (68.0)	23.8 (74.9)	25.8 (78.5)	25.9 (78.7)	22.0 (71.6)	15.9 (60.9)	10.3 (50.6)	5.1 (41.2)	15.4 (59.8)
Temp Change (1978-1987 vs. 1998-2007)	3.9 (7.0)	2.4 (4.4)	1.9 (3.5)	2.1 (3.7)	1.5 (2.7)	1.0 (1.8)	0.8 (1.5)	1.6 (2.9)	1.1 (2.0)	1.5 (2.7)	1.2 (2.2)	1.4 (2.6)	1.7 (3.1)
2007 Avg	5.0 (41.0)	3.2 (37.7)	13.9 (57.0)	13.4 (56.1)	20.9 (69.7)	24.4 (76.0)	25.1 (77.1)	28.7 (87.7)	23.6 (74.4)	17.9 (64.2)	9.4 (48.9)	7.5 (45.5)	16.1 (60.9)
Precipitation, mm (in.)													
1978-1987 Avg	103.4 (4.07)	101.4 (3.99)	112.3 (4.42)	94.5 (3.72)	123.5 (4.86)	84.4 (3.32)	137.2 (5.40)	95.0 (3.74)	85.9 (3.38)	74.4 (2.93)	120.4 (4.74)	108.5 (4.27)	1241 (48.83)
1988-1997 Avg	136.7 (5.38)	135.9 (5.35)	146.1 (5.75)	97.1 (3.82)	114.3 (4.50)	122.5 (4.82)	134.4 (5.29)	91.7 (3.61)	101.9 (4.01)	69.1 (2.72)	128.8 (5.07)	157.3 (6.19)	1436 (56.52)
1998-2007 Avg	126.8 (4.99)	122.0 (4.80)	119.7 (4.71)	151.2 (5.95)	116.6 (4.59)	117.4 (4.62)	140.5 (5.53)	61.7 (2.43)	98.3 (3.87)	63.8 (2.51)	117.6 (4.63)	108.7 (4.28)	1344 (52.91)
Precip Change (1978-1987 vs. 1998-2007)	23.4 (0.92)	20.6 (0.81)	7.4 (0.29)	56.7 (2.23)	-6.9 (-0.27)	33.0 (1.30)	3.3 (0.13)	-33.3 (-1.31)	12.4 (0.49)	-10.7 (-0.42)	-2.8 (-0.11)	0.3 (0.01)	103.7 (4.08)
2007 Avg	74.4 (2.93)	33.3 (1.31)	83.6 (3.29)	134.9 (5.31)	69.1 (2.72)	40.4 (1.59)	106.5 (4.19)	18.0 (0.71)	62.8 (2.47)	70.6 (2.78)	128.8 (5.07)	88.9 (3.50)	911.4 (35.87)
Snowfall, mm (in.)													
1978-1987 Avg	117 (4.6)	142 (5.6)	33 (1.3)	23 (0.9)	0	0	0	0	0	8 (0.3)	Trace	28 (1.1)	351 (13.8)
1988-1997 Avg	92 (3.6)	79 (3.1)	69 (2.7)	Trace	0	0	0	0	0	0	3 (0.1)	86 (3.4)	328 (12.9)
1998-2007 Avg	46 (1.8)	46 (1.8)	Trace	Trace	0	0	0	0	0	0	Trace	18 (0.7)	109 (4.3)
Precip Change (1978-1987 vs. 1998-2007)	-71 (-2.8)	-96 (-3.8)	-33 (-1.3)	-23 (-0.9)	0	0	0	0	0	-8 (-0.3)	0	-10 (0.4)	-242 (-9.5)
2007 Avg	Trace	25.4 (1.0)	Trace	Trace	0	0	0	0	0	0	0	Trace	25 (1.0)

Table C.3. Hourly freeze data for Oak Ridge, Tennessee, 1985–2007
 Number of hours at or below a given temperature (°C)^a

Year	January				February				March			April		May		October			November				December				Annual			
	≤0	<-5	<-10	<-15	≤0	<-5	<-10	<-15	≤0	<-5	<-10	≤0	<-5	≤0	<-5	≤0	<-5	<-10	≤0	<-5	<-10	<-15	≤0	<-5	<-10	<-15	≤0	<-5	<-10	<-15
1985	467	195	103	39	331	127	26	0	105	6	0	43	3	0	0	0	0	22	0	0	431	201	66	2	1399	532	195	41		
1986	308	125	38	10	161	29	3	0	124	28	0	17	0	0	0	0	0	32	10	0	232	34	0	0	874	226	41	10		
1987	302	53	7	0	111	19	3	0	95	0	0	55	4	0	0	36	0	103	18	0	151	16	0	0	853	110	10	0		
1988	385	182	43	0	294	102	19	0	97	9	0	6	0	0	0	45	0	62	3	0	301	55	0	0	1190	351	62	0		
1989	163	27	0	0	190	66	10	0	35	0	0	18	0	3	0	7	0	125	14	0	421	188	71	30	962	295	81	30		
1990	142	13	0	0	115	5	0	0	35	0	0	35	0	0	0	19	0	62	1	0	172	43	5	0	580	62	5	0		
1991	186	44	0	0	158	47	15	0	49	0	0	0	0	0	0	4	0	148	16	0	192	38	0	0	737	145	15	0		
1992	230	65	8	0	116	22	0	0	116	4	0	27	2	0	0	7	0	100	0	0	166	9	0	0	762	102	8	0		
1993	125	11	0	0	245	47	8	0	124	32	9	3	0	0	0	0	0	152	2	0	223	44	0	0	872	136	17	0		
1994	337	191	85	26	196	46	3	0	66	0	0	18	0	0	0	0	0	53	1	0	142	0	0	0	812	238	88	26		
1995	240	45	6	0	217	84	18	0	37	0	0	0	0	0	0	0	0	142	3	0	288	84	10	0	924	216	34	0		
1996	301	91	0	0	225	110	62	27	182	49	6	23	0	0	0	3	0	101	0	0	194	40	4	0	1029	290	72	27		
1997	254	101	24	0	67	0	0	0	25	0	0	6	0	0	0	6	0	96	10	0	232	14	0	0	686	125	24	0		
1998	97	10	7	0	25	0	0	0	74	20	0	0	0	0	0	0	0	38	0	0	132	4	0	0	366	34	7	0		
1999	181	68	0	0	113	14	0	0	62	0	0	0	0	0	0	4	0	41	0	0	177	23	0	0	578	105	0	0		
2000	273	62	5	0	127	30	0	0	18	0	0	8	0	0	0	11	0	94	11	0	345	124	7	0	876	227	12	0		
2001	281	60	5	0	79	9	0	0	53	0	0	2	0	0	0	18	0	28	0	0	137	35	0	0	598	104	5	0		
2002	185	28	0	0	121	16	0	0	91	17	0	2	0	0	0	0	0	41	0	0	82	6	0	0	522	67	0	0		
2003	345	123	26	0	117	12	0	0	19	0	0	0	0	0	0	0	0	37	0	0	102	9	0	0	620	144	26	0		
2004	285	50	2	0	76	0	0	0	18	0	0	0	0	0	0	0	0	9	0	0	247	41	4	0	635	91	6	0		
2005	151	65	6	0	52	1	0	0	81	1	0	0	0	0	0	1	0	55	0	0	176	28	0	0	516	95	6	0		
2006	70	0	0	0	169	19	0	0	44	0	0	0	0	0	0	15	0	37	0	0	126	41	1	0	461	60	1	0		
2007	189	30	5	0	283	70	0	0	29	0	0	32	0	0	0	0	0	60	0	0	83	8	0	0	673	111	5	0		
Avg.	239	71	16	3	156	38	7	1	69	7	1	13	0	0	0	8	0	71	4	0	207	47	7	1	762	168	31	6		

^aSource: 1985–2007 National Oceanic and Atmospheric Administration Atmospheric Turbulence and Diffusion Division KOQT Station, Automated Surface Observing System.

