Abstract Submitted
for the DNP02 Meeting of
The American Physical Society

Sorting Category: 5 (Experimental)

Breakup of ^{17}F by ^{208}Pb near the Coulomb barrier

J.F. LIANG, J.R. BEENE, A. GALINDO-URIBARRI, J. GOMEZ DEL CAMPO, C.J. GROSS, P.A. HAUSLADEN, Y. LAROCHELLE, P.E. MUELLER, D. SHAPIRA, D.W. STRACENER, R.L. VARNER, Physics Division, Oak Ridge National Laboratory[1], H. ESBENSEN, Physics Division, Argonne National Laboratory[2], J.D. BIERMAN, Physics Department AD-51, Gonzaga University — Angular distributions of ^{16}O from 98 and 120 MeV ^{17}F on ^{208}Pb were measured with a ΔE-E telescope which consists of a 5×5 cm^2 Si detector and a double-sided Si strip detector. A peak was observed forward of the grazing angle. Full dynamical calculations show that the measured angular distributions are dominated by the stripping breakup reaction.

[1]Oak Ridge National Laboratory is supported by the U.S. Department of Energy under contract No. DE-AC05-00OR22725 with UT-Battelle, LLC.
[2]Supported by the U.S. Department of Energy, Nuclear Physics Division, under contract No. W-31-109-ENG-38.

Junjien Liang
liang@mail.phy.ornl.gov
Physics Division, Oak Ridge National Laboratory
Member ID: LI042361

[X] Prefer Oral Session
[ ] Prefer Poster Session

Special instructions: primary

Date submitted: June 28, 2002 Electronic form version 1.4