

A spectroscopic view of electron-phonon coupling at metal surfaces

Junren Shi², S.-J Tang¹, Biao Wu^{2,3}, P.T. Sprunger⁴, W.L. Yang^{5,6}, V. Brouet^{5,6}, X.J. Zhou⁵, Z. Hussain⁶, Z.-X. Shen⁵, Zhenyu Zhang^{1,2}, and E.W. Plummer^{1,2}

¹Department of Physics and Astronomy, University of Tennessee, Knoxville, TN 37996;

²Condensed Mater Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831;

³Department of Physics, University of Texas, Austin, TX 78712; ⁴Department of Physics and

Astronomy, Louisiana State University, Baton Rouge, LA 70803; ⁵Department of Physics,

Applied Physics and Stanford Synchrotron Radiation Laboratory, Stanford University, Stanford,

CA 94305; ⁶Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, CA

94720.

eplummer@utk.edu

Electron-phonon interaction in metals is a very old subject. Göran Grimvall in the preface to his classic book on this subject pointed out that 1000 papers had been published on this subject between 1960 and 1980 [1]. What can be new and exciting in such an old and mature field? The answer is *surfaces and surface science* techniques. Recent advances in experimental techniques coupled with ever-growing theoretical capabilities promise to create a renaissance in this subject. High-resolution angular-resolved photoemission is producing direct images of the distortion of the two-dimensional surface state bands near the Fermi energy caused by electron-phonon coupling (EPC). First-principles calculations of the EPC for surface states are appearing in the literature [2] which not only explain the origin of the EPC-induced band distortions but also produce exquisite pictures of the Eliashberg function $\alpha^2F(\omega)$ [1]. The Eliashberg function is the product of a coupling constant times the phonon density of states and is at the heart of any theory of EPC [1]. A recent theoretical advance that will be described in this talk will allow experimentalists to extract the Eliashberg function directly from the high-resolution photoemission data [4]. These developments mark the beginning of a new era or a renaissance in the elucidation of many-body effects in reduced dimensionality.

In the 1960s the most definitive signature for determining the mechanism in conventional superconductors was the measurement of the single-particle tunneling I-V characteristic and the concomitant inversion procedure to display the Eliashberg function. Today EPC is intimately associated with the functionality of complex materials and its role in high- T_c superconductivity is being actively discussed. Research on metal surfaces can produce in unprecedented detail a picture of electron-phonon coupling. The Eliashberg function can be measured as a function of both E and \mathbf{k} and related directly to measured and calculated surface phonon dispersion curves. Several recent examples will be used to illustrate the detailed spectroscopic information that can be extracted from the high-resolution angle-resolved photoemission data.

These investigations on surfaces of relatively simple metals can be the platform for understanding functionality in complex materials associated with the coupling between charge and the lattice. EPC as displayed by surface states on $Be(0001)$ and $Be(1010)$ will be used to illustrate the power of the direct extraction technique [3] and the physical mechanisms responsible for the large EPC at the surfaces. Finally, recent data on the under doped cuprate High- T_c superconductor, $La_{2-x}Sr_xCuO_4$, in the normal state will be described.

References:

1. G. Grimvall, in *The Electron-Phonon Interaction in Metals, Selected Topics in Solid State Physics*, edited by E. Wohlfarth (North-Holland, New York, 1981).
2. A. Eiguren, S. de Gironcoli, E. V. Chulkov, P. M. Echenique, and E. Tosatti, *Phys. Rev. Lett.* **91**, 166803 (2003).
3. Junren Shi, S.-J. Tang, Biao Wu, P. T. Sprunger, W. L. Yang, V. Brouet, X. J. Zhou, Z. Hussain, Z.-X. Shen, Zhenyu Zhang, and E. W. Plummer, *Phys. Rev. Letters* to be published.