

# The Impact of Nuclear Reaction Rate Uncertainties on Evolutionary Studies of the Nova Outburst

W. Raphael Hix<sup>†§</sup>, Michael S. Smith<sup>§</sup>, Anthony Mezzacappa<sup>§</sup>,  
Sumner Starrfield<sup>\*</sup>, Donald L. Smith<sup>‡</sup>

<sup>†</sup>*Department of Physics & Astronomy, University of Tennessee, Knoxville, TN 37996-1200*

<sup>§</sup>*Physics Division, Oak Ridge National Laboratory<sup>1</sup> Oak Ridge, TN 37831-6354*

<sup>\*</sup>*Department of Physics & Astronomy, Arizona State University, Tempe, AZ 85287-1504*

<sup>‡</sup>*Technology Development Division, Argonne National Laboratory, Argonne, IL 60439*

**Abstract.** The observable consequences of a nova outburst depend sensitively on the details of the thermonuclear runaway which initiates the outburst. One of the more important sources of uncertainty is the nuclear reaction data used as input for the evolutionary calculations. A recent paper by Starrfield, Truran, Wiescher, & Sparks [1] has demonstrated that changes in the reaction rate library used within a nova simulation have significant effects, not just on the production of individual isotopes (which can change by an order of magnitude), but on global observables such as the peak luminosity and the amount of mass ejected. We will present preliminary results of systematic analyses of the impact of reaction rate uncertainties on nova nucleosynthesis.

## I NUCLEAR UNCERTAINTIES IN NOVAE

Observations of nova outbursts have revealed an elemental composition that differs markedly from solar. Theoretical studies indicate that these differences are caused by the combination of convection with explosive hydrogen burning which results in a unique nucleosynthesis that is rich in odd-numbered nuclei such as  $^{15}\text{N}$ ,  $^{17}\text{O}$  and  $^{13}\text{C}$ . These nuclei are difficult to form in other astrophysical events. Many of the proton-rich nuclei produced in nova outbursts are radioactive, offering the possibility of direct observation with  $\gamma$ -ray instruments. Potentially important  $\gamma$ -ray producers include  $^{26}\text{Al}$ ,  $^{22}\text{Na}$ ,  $^7\text{Be}$  and  $^{18}\text{F}$ .

The observable consequences of a nova outburst depend sensitively on the details of the thermonuclear runaway which initiates the outburst. One of the more important sources of uncertainty is the nuclear reaction data used as input for the

---


<sup>1)</sup> Research at the Oak Ridge National Laboratory is supported by the U.S. Department of Energy under contract DE-AC05-96OR22464 with Lockheed Martin Energy Research Corp.

evolutionary calculations [1]. A number of features conspire to magnify the effects of nuclear uncertainties on nova nucleosynthesis. Many reactions of relevance to novae involve unstable proton-rich nuclei, making experimental rate determinations difficult. For hydrodynamic conditions typical of novae, many rates depend critically on the properties of a few individual resonances, resulting in wide variation between different rate determinations. Statistical model (Hauser-Feshbach) calculations, which are employed with great success for a large number of reactions [2], are unreliable for rates dominated by individual resonances. The similarity of the nuclear burning and convective timescales results in nuclear burning in novae which is far from the steady state which typifies quiescent burning.

## II MONTE CARLO ESTIMATES OF UNCERTAINTIES

Though analysis of the impact of variations in the rates of a few individual re-

,




**FIGURE 1.** Final Abundances (elapsed time =  $5.2 \times 10^5$  sec after peak).

where  $\alpha$  and  $\beta$  are the (logarithmic) mean and standard deviation. For small uncertainties ( $< 20\%$ ), the difference between the log-normal distribution and the normal (Gaussian) distribution is small. However, for uncertainties of larger sizes such as those encountered in this problem, the difference is important. For this preliminary analysis, we have chosen to assign uncertainties of  $\sim 50\%$  ( $\beta = \ln(1.5)$ ) both to rates calculated by Hauser-Feshbach methods and also to rates whose measurement require radioactive ion beams. For all other rates we assign  $\beta = \ln(1.2)$ . Figure 2 plots the resulting abundance distributions for two representative nuclei. Fig. 2 also demonstrates the differences between normal and log-normal distributions for widths of these sizes. These are very conservative uncertainties; relatively few reactions, especially among unstable nuclei, have measurement uncertainties this small.

### III RESULTS

As evidenced by the error bars in Fig. 1, the impact of even our conservatively chosen variations in reaction rates on the nucleosynthesis is large. While broader conclusions will require analysis of additional hydrodynamic trajectories, a number of interesting points can be made from the analysis of this single trajectory. The impact on the rate of energy production is small. At the 90% confidence level,



**FIGURE 2.** The histograms showing deviations in abundance from the mean. The dotted curves are normal distributions with the same (arithmetic) mean and standard deviation as the Monte Carlo distribution. The dashed curves are log-normal distributions with (logarithmic) mean and standard deviation from the Monte Carlo. Y Center, Y Upper, and Y Lower correspond, respectively, to the central abundance and the upper and lower error bars from Fig. 1.

variations in the amount of hydrogen consumed represent  $\sim 10\%$  variation in the thermonuclear energy released. For the most abundant metals (those which represent more than 1% of the mass),  $2\sigma$  variations by factors of 1.1 to 1.4 are common, with some of these nuclei showing  $2\sigma$  variations as large as a factor of 2, for example,  $^{15}\text{N}$  ( $2.1\times$ ) and  $^{30}\text{Si}$  ( $2.3\times$ ). For the  $\gamma$ -ray source nuclei  $^{22}\text{Na}$  and  $^{26}\text{Al}$ , the 90% confidence interval includes variations of nearly a factor of 2, representing almost a factor of 4 difference in the distance from which novae may be observed by  $\gamma$ -ray telescopes. For  $^7\text{Be}$ , the 90% confidence interval spans more than 2 orders of magnitude.

Such large uncertainties in the nucleosynthesis, resulting from poorly known nuclear reaction rates, constrain our ability to make detailed comparisons between theoretical models for the nova outburst and astrophysical observations to a degree which is often ignored. Improved knowledge of these uncertain rates, both experimental and theoretical, is necessary to provide tight constraints on the nova outburst from its nucleosynthetic products.

## REFERENCES

1. S. Starrfield, J. W. Truran, M. C. Wiescher, and W. M. Sparks, *MNRAS* **296**, 502 (1998).
2. T. Rauscher, F.-K. Thielemann, and K.-L. Kratz, *Nucl. Phys. A* **621**, 331 (1997).
3. J. José, A. Coc, and M. Hernanz, *ApJ* **520**, 347 (1999).
4. W. Hillebrandt and F.-K. Thielemann, *ApJ* **255**, 617 (1982).
5. M. Wiescher, J. Gorres, F.-K. Thielemann, and H. Ritter, *A&A* **160**, 56 (1986).
6. M. Politano *et al.*, *ApJ* **448**, 807 (1995).
7. M. S. Smith, L. H. Kawano, and R. A. Malaney, *ApJS* **85**, 219 (1993).
8. D. L. Smith, *Probability, Statistics, and Data Uncertainties in Nuclear Science and Technology* (LaGrange Park: Am. Nuc. Soc., 1991).